

ISBN 82-553-0488-6

Mathematics
May 27

No.9
1982

DUALITY FOR RATIONAL NORMAL SCROLLS

by

Ragni Piene	and	Gianni Sacchiero
Institute of Math.		Istituto Matematico
University of Oslo		Università di Ferrara

DUALITY FOR RATIONAL NORMAL SCROLLS

by

Ragni Piene and Gianni Sacchiero

1. Introduction

Given a skew curve $C \subset \mathbb{P}^n$, its strict dual curve $C^* \subset \mathbb{P}^n$ is defined as the closure of the set of hyperplanes that are osculating to C of order n at smooth points of C . The strict dual of C^* is C , and there is a natural duality between the various osculating developables of C and those of C^* [P1, §5]. If $C \subset \mathbb{P}^n$ is a rational normal curve, then so is $C^* \subset \mathbb{P}^n$; we say that C is self-dual.

In this paper we shall consider rational normal scrolls $X \subset \mathbb{P}^N$ of dimension r , and study their higher order dual varieties $X_m^V \subset \mathbb{P}^N$, in particular the strict dual variety $X^* \subset \mathbb{P}^N$. We show that only the generic scrolls (in the sense of Ghione [G]) satisfy the biduality $X^{**} = X$, and that, among these, only the balanced scrolls are self-dual. By defining the "restriction" of X^* to sub-scrolls of X , the nature of the self-duality for balanced scrolls appears even clearer.

Similar considerations could be applied to non-rational scrolls and to non-normal scrolls (e.g., projections of normal ones). Except for statements like: "if there exists a directrix of low degree, then the dual varieties are contained in a lower-dimensional space, and so biduality does not hold", it is not so simple, however, to state general results in those cases.

2. Rational normal scrolls

Fix integers $0 < d_1 \leq \dots \leq d_r$, with $d_r > 0$, and consider the rank r bundle

$$F = \bigoplus_{i=1}^r \mathcal{O}_{\mathbb{P}^1}(d_i)$$

on $\mathbb{P}^1 = \mathbb{P}_k^1$, where k is an algebraically closed field of characteristic 0. The complete linear system associated with the line bundle $\mathcal{O}_{\mathbb{P}(F)}(1)$ defines a morphism

$$\phi : \mathbb{P}(F) \rightarrow \mathbb{P}(V) \simeq \mathbb{P}^N$$

where $V = \bigoplus_{i=1}^r V_i$, $V_i = H^0(\mathcal{O}_{\mathbb{P}^1}(d_i))$, and $N = \sum_{i=1}^r (d_i+1)-1 = \sum_{i=1}^r d_i + r - 1$. Set $X = \phi(\mathbb{P}(F))$; then we have $\deg X = \sum_{i=1}^r d_i$. We shall say that X is a variety of type (d_1, \dots, d_r) .

If $d_1 > 0$ holds, $\mathcal{O}_{\mathbb{P}(F)}(1)$ is very ample and ϕ is an embedding. Then we call X a rational normal scroll of dimension r . If $d_1 = \dots = d_s = 0 < d_{s+1}$ for some s , $1 \leq s < r$, then X is a cone, with a \mathbb{P}^{s-1} as vertex, over a rational normal scroll of type (d_{s+1}, \dots, d_r) .

Classically (C. Segre [S1],[S2]) these varieties were constructed in the following way: Take linear subspaces $\mathbb{P}^{d_i} \subset \mathbb{P}^N$, $i = 1, \dots, r$, such that no \mathbb{P}^{d_i} intersects the space spanned by the other \mathbb{P}^{d_j} 's. Consider now the d_i -uple embedding $\phi_i: \mathbb{P}^1 \rightarrow \mathbb{P}^{d_i}$; thus we have in each \mathbb{P}^{d_i} a rational normal curve $D_i = \phi_i(\mathbb{P}^1)$. (If $d_i = 0$, D_i is just the point \mathbb{P}^0 .) The scroll X is then the r -dimensional variety swept out by the $(r-1)$ -dimensional subspaces $X_t = [\phi_1(t), \dots, \phi_r(t)]$, as t varies in \mathbb{P}^1 . We call the X_t 's the generators of X .

Finally, we shall also use the notation

$$X = (D_1, \dots, D_r) .$$

3. Higher order dual varieties

Recall the definition [P2] of higher order dual varieties of a given variety $X \subset \mathbb{P}(V) \simeq \mathbb{P}^n$: For each $m > 0$, consider the homomorphism

$$a^m: V_X \rightarrow P_X^m(1),$$

where $P_X^m(1)$ is the sheaf of principal parts of order m of $O_X(1)$. Let $s = s(m)$ denote the generic rank of a^m . If $s \leq n$, then almost all points of X has a well-defined $(s-1)$ -dimensional osculating space of order m . The m -th dual variety $X_m^V \subset \mathbb{P}(V^V) \simeq \mathbb{P}^n$ of X is then defined as the closure of the set of hyperplanes containing an m -th order osculating $(s-1)$ -space.

Set $\bar{s} = \max\{s(m) \mid s(m) \leq n, m \geq 0\}$, and $\bar{m} = \min\{m \mid s(m) = \bar{s}\}$. Then we call $X^* = X_{\bar{m}}^V$ the strict dual variety of X . This definition, being slightly more general than the one given in [P2], permits us to define the strict dual of any variety $X \subset \mathbb{P}^n$.

Assume now that $X \subset \mathbb{P}^n$ is a variety of type (d_1, \dots, d_r) . The coordinates $x_0^i, \dots, x_{d_i}^i$ on \mathbb{P}^{d_i} give coordinates $x_0^1, \dots, x_{d_1}^1, x_0^2, \dots, x_{d_2}^2, \dots, x_0^r, \dots, x_{d_r}^r$ on \mathbb{P}^N . If t is an affine parameter on \mathbb{P}^1 , then X is given parametrically at points $x \in X$ around the generator X_t by

$$x_j^i = x_j^i(\lambda_1, \dots, \lambda_r; t) = \lambda_i t^j$$

($i = 1, \dots, r$; $j = 0, 1, \dots, d_i$), where $(\lambda_1, \dots, \lambda_r) \in \mathbb{P}^{r-1}$.

Since X has dimension r , the sheaf $P_X^m(1)$ has generic rank $\binom{t+m}{m}$. (If X is smooth, i.e., if $d_1 > 0$, then $P_X^m(1)$ is a bundle with this rank.) Since X is ruled (the parametrization is linear in the λ_i 's), the generic rank of

$$a^m: V_X \rightarrow P_X^m(1)$$

is, however, always $\leq rm+1$, and clearly equality holds if and only if $m \leq d_1$. Assume, say, that $\lambda_r \neq 0$. Then the image of a^m is

given by the matrix

$$A_m = \begin{pmatrix} (h_1) \\ x_{\lambda_1} \\ \vdots \\ (h_{r-1}) \\ x_{\lambda_{r-1}} \\ (h_r) \\ x \end{pmatrix}$$

where $h_i = 0, 1, \dots, \min\{d_i, m-1\}$ for $1 \leq i \leq r-1$ and $h_r = 0, 1, \dots, \min\{d_r, m\}$, and where we set

$$x = (x_j^i), \quad x^{(h)} = \frac{\partial h_x}{\partial t^h}, \quad x_{\lambda_i}^{(h)} = \frac{\partial x^{(h)}}{\partial \lambda_i}.$$

We shall now rewrite this matrix A_m which determines x_m^v , using the following notation:

Let $D \subset \mathbb{P}^d$ be the rational normal curve $x_i = t^i$, $i = 0, 1, \dots, d$. For each $m \geq 0$, set

$$M_d^m = \begin{pmatrix} x \\ x' \\ \vdots \\ x^{(m)} \end{pmatrix}.$$

If $m \leq d$, then

$$M_d^m = \begin{pmatrix} 1 & t & \dots & \dots & \dots & t^d \\ 0 & 1 & & & & \vdots \\ \vdots & \vdots & \ddots & & & \vdots \\ 0 & \dots & \dots & 0 & 1 & \dots & \binom{d}{m} t^{d-m} \end{pmatrix}$$

and if $m > d$, then

$$M_d^m = \begin{pmatrix} M_d^d \\ 0 \end{pmatrix}$$

We shall write

$$\bar{M}_d^m = \begin{cases} M_d^m & \text{if } m \leq d \\ M_d^d & \text{if } m > d \end{cases}$$

The matrix \bar{M}_d^m corresponds to the map

$$a^m : V_D \rightarrow P_D^m(1)$$

giving the m -th order osculating spaces (or the m -th osculating developable, or the m -th dual variety) of $D \subset \mathbb{P}^d$. Note that a^d is an isomorphism.

Returning to $X \subset \mathbb{P}^N$, the matrix A_m can now be written as

$$A_m = \begin{pmatrix} \bar{M}_{d_1}^{m-1} & & & & & & 0 \\ & \bar{M}_{d_2}^{m-1} & & & & & \\ & & \ddots & & & & \\ & & & \ddots & & & \\ & 0 & & & \ddots & & \\ & & & & & \bar{M}_{d_{r-1}}^{m-1} & \\ \lambda_1 M_{d_1}^m & \cdot & \cdot & \cdot & \cdot & \lambda_{r-1} M_{d_{r-1}}^m & M_{d_r}^m \end{pmatrix}$$

for $1 \leq m \leq \bar{m}$. It follows that the m -th order osculating space to X at a point $x \in D_r \cap X_t$, is equal to the space spanned by the $(m-1)$ -th order osculating space to D_i at $x_i = D_i \cap X_t$, $i = 1, \dots, r-1$, and the m -th order osculating space to D_r at x . (If $m-1 > d_i$, then any $(m-1)$ -th order osculating space to D_i is the whole space $[D_i]$.)

Similar considerations apply to the cases $\lambda_i \neq 0$. In particular, we observe that the intersection of the m -th order osculating spaces to X at points of a generator X_t contains the space spanned by the $(m-1)$ -th order osculating spaces to the directrices D_i at the points $x_i = D_i \cap X_t$.

Before stating the general results, we shall give some, hopefully illuminating, examples.

4. Examples

We shall first consider some surfaces.

Example 1. $r = 2$, $\deg X = 4$, $N = 5$.

There are three possible types:

- a) $d_1 = 0$, $d_2 = 4$
- b) $d_1 = 1$, $d_2 = 3$
- c) $d_1 = d_2 = 2$.

Case a): For $1 \leq m \leq \bar{m} = 3$, A_m becomes

$$\begin{pmatrix} M_0^{m-1} & 0 \\ \lambda_1 M_0^m & M_4^m \end{pmatrix}.$$

The strict dual variety $X^* = X_3^V$ is given by

$$\begin{pmatrix} 1 & 0 \\ 0 & M_4^3 \end{pmatrix}.$$

The points of X^* are thus the 4-spaces spanned by an osculating 3-space to D_2 and the point D_1 . In other words,

$$X = (D_2)_3^V \cap [D_1]^V.$$

Note that $(D_2)_3^V$ is a cone with vertex $[D_2]^V$. So X^* is a rational normal curve of degree 4 in the hyperplane $[D_1]^V \subset \mathbb{P}^5$.

Moreover, we observe that

$$X_2^V = (D_2)_2^V \cap [D_1]^V$$

is the tangent developable of $X^* \subset [D_1]^V$, and that

$$X^V = X_1^V = (D_2)_1^V \cap [D_1]^V$$

is the osculating developable of $X^* \subset [D_1]^V$ ([P1], 5.2).

Case b): For $1 \leq m \leq \bar{m} = 2$, A_m becomes

$$\begin{pmatrix} M_1^{m-1} & 0 \\ \lambda_1 M_1^m & M_3^m \end{pmatrix},$$

so $X^* = X_2^V$ is given by the matrix (equivalent to A_m)

$$\begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ - & - & - & - \\ 0 & & 1 & M_3^2 \end{pmatrix}$$

Hence $X^* = (D_2)_2^V \cap [D_1]^V$ is a rational normal curve of degree 3.

Case c): For $1 \leq m \leq \bar{m} = 2$, A_m is

$$\begin{pmatrix} M_2^{m-1} & 0 \\ \lambda_1 M_2^m & M_2^m \end{pmatrix}$$

At this point it is convenient to introduce the "restriction of $X^* = X_2^V$ to D_i ":

$X^*|_{D_i}$ = the points of X^* corresponding to points on D_i .

Then $X^*|_{D_2}$ is obtained by setting $\lambda_1 = 0$, hence

$$X^*|_{D_2} = (D_1)_1^V \cap [D_2]^V.$$

Similarly,

$$X^*|_{D_1} = (D_2)_1^V \cap [D_1]^V.$$

We shall call $D_i^* = (D_i)_{d_{i-1}}^V \cap [D_{3-i}]^V$, $i = 1, 2$, the strict dual curve of D_i (with respect to D_{3-i}). (Note that there is a self-duality for rational normal curves between $D_i \subset [D_i]$ and $D_i^* \subset [D_i^*] = [D_{3-i}]^V$.)

With this notation we have

$$X^*|_{D_i} = D_{3-i}^*, \quad i = 1, 2,$$

and

$$X^* = (D_1^*, D_2^*).$$

Hence X^* is again of type $(2, 2)$, and so X is self-dual.

Example 2. $r = 2$, $\deg X = 5$, $N = 6$.

There are three possible types:

- a) $d_1 = 0$, $d_2 = 5$
- b) $d_1 = 1$, $d_2 = 4$
- c) $d_1 = 2$, $d_2 = 3$

The cases a) and b) are similar to the case a) and b) of Ex.1. Here we have $\bar{m} = d_2 - 1$ and $\dim X^* = 1$. In fact, $X^* = D_2^* \subset [D_1]^V$.

c): $\bar{m} = 2$.

$$A_2 = \begin{pmatrix} M_2^1 & 0 \\ \lambda_1 M_2^2 & M_2^3 \end{pmatrix}$$

$X^*|D_1$ is given by

$$\begin{pmatrix} M_2^1 & 0 \\ 0 & M_2^3 \end{pmatrix}$$

which is a (7×5) matrix, and hence $X^*|D_1 = (D_1^*, D_2^*)$ is a ruled surface. Now $X^*|D_2$ is given by

$$\begin{pmatrix} 0 & M_2^1 \\ M_2^2 & 0 \end{pmatrix}$$

so $X^*|D_2 = (D_2^*)_1^V \cap [D_1]^V$ is equal to the tangent developable $S^1(D_2^*)$ of D_2^* in $[D_1]^V$ ([P1], §5).

Hence X^* is determined by the ruled surface (D_1^*, D_2^*) and the developable surface $S^1(D_2^*)$, which also contains D_2^* .

In other words, X^* is determined by $D_1^* \subset [D_2]^V$ and $S^1(D_2^*) \subset [D_1]^V$. We shall write, with abuse of notation, $X^* = (D_1^*, S^1(D_2^*))$. Hence X^* is a rational (non-normal) 3-dimensional scroll of type $(2, 2, 2)$ (in the obvious sense) because $S^1(D_2^*)$ is of type $(2, 2)$.

Moreover, $X^{**} = (X^*)^{\vee}_2 = X$; this can be seen by the fact that $X^{**}|D_1^* = D_2$ and $X^{**}|S^1(D_2^*) = D_1$.

The preceding examples are typical for surfaces, in the following sense: If $X = (D_1, D_2)$, then for $m \leq \bar{m}$,

$$\dim X_m^{\vee} = \begin{cases} N+1-2m & m \leq d_1 \\ d_2-m & d_1 < m < d_2 \end{cases}$$

For the strict dual $X^* = X_{\bar{m}}^{\vee}$, there are three possible cases:

- a) $d_1 = d_2 = d$: $\bar{m} = d$, $X^* = (D_1^*, D_2^*)$.
- b) $d_1 = d$, $d_2 = d+1$: $\bar{m} = d$, $X^* = (D_1^*, S^1(D_2^*))$ is a 3-dimensional (non-normal) scroll of type (d, d, d) .
- c) $d_1+2 \leq d_2$: $\bar{m} = d_2-1$, $X^* = D_2^*$. Moreover, X_m^{\vee} , for $d_1+1 \leq m \leq d_2-1$, is equal to $S^{d_2-1-m}(D_2^*) \subset [D_1]^{\vee}$.

Hence X is self-dual only in case a), and only in the cases a) and b) does biduality ($X = X^{**}$) hold.

Example 3. $r = 3$, $\deg X = 6$, $N = 8$.

There are 7 possible cases for the type (d_1, d_2, d_3) of X :

- a) $(0, 0, 6)$
 - b) $(0, 1, 5)$
 - c) $(0, 2, 4)$
 - d) $(1, 1, 4)$
 - e) $(1, 2, 3)$
 - f) $(0, 3, 3)$
 - g) $(2, 2, 2)$
- $\left. \begin{matrix} a) \\ b) \\ c) \\ d) \end{matrix} \right\} \bar{m} = d_3-1, \quad X^* = D_3^* \subset [D_1, D_2]^{\vee}$
 $\left. \begin{matrix} e) \\ f) \end{matrix} \right\} \bar{m} = d_2, \quad X^* = (D_2, D_3)^* \subset [D_1]^{\vee}$
 $g) \quad \bar{m} = 2, \quad X^* = (D_1^*, D_2^*, D_3^*).$

In the first four cases, the matrix A_m is equivalent to (i.e., defines the same variety as) the matrix

$$\begin{pmatrix} M_{d_1}^{d_1} & 0 & 0 \\ 0 & M_{d_2}^{d_2} & 0 \\ 0 & 0 & M_{d_3}^{d_3-1} \end{pmatrix}$$

In the cases e) and f), the matrix $A_{\overline{m}} = A_{d_2}$ is equivalent to

$$\begin{pmatrix} M_{d_1}^{d_1} & 0 & 0 \\ 0 & M_{d_2}^{d_2-1} & 0 \\ 0 & \lambda_2 M_{d_2}^{d_2} & M_{d_3}^{d_2} \end{pmatrix}$$

The conclusions follow, by considerations as in the previous examples.

Consider now the case g). The matrix $A_{\overline{m}} = A_2$ becomes

$$\begin{pmatrix} M_2^1 & 0 & 0 \\ 0 & M_2^1 & 0 \\ \lambda_1 M_2^2 & \lambda_2 M_2^2 & M_2^2 \end{pmatrix}$$

As in Ex.1 c) we consider the restriction $X^*|D_3$. This is the variety defined by setting $\lambda_1 = \lambda_2 = 0$ in A_2 . Hence

$$X^*|D_3 = (D_1^*, D_2^*) \subset [D_3]^V.$$

In the same way, we obtain

$$X^*|D_1 = (D_2^*, D_3^*) \subset [D_1]^V$$

$$X^*|D_2 = (D_1^*, D_3^*) \subset [D_2]^V$$

This shows that $X^* = (D_1^*, D_2^*, D_3^*)$, and hence that X is self-dual.

Since the restriction of X^* to a directrix D_i is the ruled surface defined by the strict duals of the other D_j 's, it is natural to ask whether the converse is true.

That is, we want to give a natural definition of the restriction $X^*|(D_2, D_3)$, so that it becomes equal to D_1^* .

To do this, we take for $X^*|(D_2, D_3)$ the points in \mathbb{P}^N corresponding to the hyperplanes that are 2-osculating to X at every point of some generators of (D_2, D_3) . This means that we should look at the matrix

$$\begin{pmatrix} M_2^1 & 0 & 0 \\ 0 & M_2^2 & 0 \\ 0 & \lambda_2 M_2^2 & M_2^2 \end{pmatrix},$$

which gives $X^*|(D_2, D_3) = D_1^*$.

Example 4. $r = 3$, $\deg X = 7$, $N = 9$.

The following types are possible:

- a) $(0,0,7)$
 - b) $(0,1,6)$
 - c) $(0,2,5)$
 - d) $(1,1,5)$
 - e) $(1,2,4)$
 - f) $(0,3,4)$
 - g) $(1,3,3)$
 - h) $(2,2,3)$
- $\left. \begin{array}{l} \text{c) } (0,2,5) \\ \text{d) } (1,1,5) \\ \text{e) } (1,2,4) \end{array} \right\} \bar{m} = d_3 - 1, X^* = D_3^* \subset [D_1, D_2]^\vee$
 $\left. \begin{array}{l} \text{f) } (0,3,4) \\ \text{g) } (1,3,3) \end{array} \right\} \bar{m} = d_2, X^* = (D_2, D_3)^* \subset [D_1]^\vee$
 $\text{h) } (2,2,3) \quad \bar{m} = 2, X^* = (D_1^*, D_2^*, S^1(D_3^*)) \text{ is of dimension } 4.$

The 7 first cases are as before; the only case that presents something new is case h). By looking at the matrix A_2 in this case, we obtain the following:

$$\begin{aligned}
 X^*|_{D_1} &= (D_2^*, S^1(D_3^*)) \subset [D_1]^\vee \\
 X^*|_{D_2} &= (D_1^*, S^1(D_3^*)) \subset [D_2]^\vee \\
 X^*|_{D_3} &= (D_1^*, D_2^*, D_3^*) \\
 X^*|(D_1, D_2) &= S^1(D_3^*) \subset [D_1, D_2]^\vee \\
 X^*|(D_1, D_3) &= (D_2^*, D_3^*) \subset [D_1]^\vee \\
 X^*|(D_2, D_3) &= (D_1^*, D_3^*) \subset [D_2]^\vee .
 \end{aligned}$$

5. Duality results

Let $X = (D_1, \dots, D_r)$ be a variety of type (d_1, \dots, d_r) . Given $(i_1, \dots, i_s) \subset (1, \dots, r)$, we denote by

$$X_{i_1, \dots, i_s} = (D_{i_1}, \dots, D_{i_s});$$

this is a variety of type (i_1, \dots, i_s) .

Set $(j_1, \dots, j_{r-s}) = (1, \dots, r) - (i_1, \dots, i_s)$.

With these notations, we give the following

DEFINITION: i) The strict dual curve D_j^* of D_j with respect to $D_{i_1}, \dots, D_{i_{r-1}}$, is the curve

$$D_j^* = (D_j)_{d_j-1}^\vee \cap [X_{i_1, \dots, i_{r-1}}]^\vee.$$

ii) The restriction of X^* to X_{i_1, \dots, i_s} is the variety $X^*|_{X_{i_1, \dots, i_s}}$ in \mathbb{P}^N consisting of hyperplanes in \mathbb{P}^N that are \bar{m} -osculating to X at every point of some generator of X_{i_1, \dots, i_s} .

PROPOSITION 1: Let $X = (D_1, \dots, D_r)$ be a variety of type (d_1, \dots, d_r) , and denote by \bar{m} the integer such that $X^* = X_{\bar{m}}^V$. For each m , $1 \leq m \leq \bar{m}$, let $i = i(m)$ denote the integer such that $0 \leq i \leq r-1$ and $d_i + 1 \leq m \leq d_{i+1}$ (for $i=0$ we set $d_0=0$). If $i \leq r-2$, then

$$\dim X_m^V = N+1-rm + \sum_{j=1}^i (m-1-d_j).$$

If $i = r-1$, then $\dim X_m^V = N-rm + \sum_{j=1}^{r-1} (m-1-d_j) = d_r - m$.

Moreover, if $i \geq 1$, we have

$$X_m^V = (X_{i+1}, \dots, r)_m^V \cap [X_1, \dots, i]^V.$$

REMARK: For $m = 1$, we obtain $\dim X_1^V \leq N+1-r$. Suppose $r \geq 2$. Then $X^V = X_1^V$ is a hypersurface if and only if $r = 2$ and $i = 0$, i.e. if and only if X is a 2-dimensional, smooth scroll.

PROPOSITION 2: With the notation of Prop. 1, we have the following description of the strict dual X^* of X .

- 1) $d_{r-1} = d_r$. Let i be the integer such that $0 \leq i < r-1$, $d_i < d_{i+1} = d_{i+2} = \dots = d_{r-1} = d_r = d$.

Then

$$X^* = X_d^V = (D_{i+1}^*, \dots, D_r^*) \subset [X_1, \dots, i]^V.$$

- 2) $d_{r-1} + 1 = d_r$. Let i be such that $0 \leq i < r-1$, $d_i < d_{i+1} = \dots = d_{r-1} = d$.

Then

$$X^* = X_d^V = (D_{i+1}^*, \dots, D_{r-1}^*, S^1(D_r^*)) \subset [X_1, \dots, i]^V.$$

m -osculating to X at all points of some linear $(r-2)$ -dimensional subspace Y_t of X_t . In fact, H is $(m-1)$ -osculating to X at all points of X_t , so $X \cap H$ contains X_t $m-1$ times, and the residual intersection is a (possibly reducible) ruled variety Y of dimension $r-1$. Hence $X \cap H$ contains $Y_t = Y \cap H$ m times, i.e., H is m -osculating to X along Y_t .

From this observation it follows that X_m^V is determined by the matrix obtained from A_m by fixing $r-2$ of the λ_i 's. That is, we fix a ruled surface $R \subset X$. If $H \in X_m^V$, then H is m -osculating to X along $Y_t \subset X_t$ (some t), hence at some point of R_t , since $R_t \cap Y_t \neq \emptyset$.

Assume therefore $\lambda_1 = \dots = \lambda_{r-2} = 0$. Then the variation in the m -th order osculating spaces along a generator is given by the variation of the matrix

$$\begin{pmatrix} M_{d_{r-1}}^{m-1} & 0 \\ \lambda_{r-1} M_{d_{r-1}}^m & M_{d_r}^m \end{pmatrix}.$$

If $m \leq d_{r-1}$ (i.e., if $i \leq r-2$), then we see that the m -osculating spaces vary with λ_{r-1} (i.e., are non-constant along R_t). Since they also vary with t , we obtain a 2-dimensional family of m -osculating spaces to X . Each such space has dimension equal to

$$rm - \sum_{j=1}^i (m-1-d_j),$$

hence the m -dual variety X_m^V has dimension

$$\dim X_m^V = N - rm + \sum_{j=1}^i (m-1-d_j) - 1 + 2 = N+1 - rm + \sum_{j=1}^i (m-1-d_j).$$

Suppose now $i = r-1$, so that $d_{r-1} < m < d_r$. Then the variation of the m -osculating spaces is given by the variation of the matrix $M_{d_r}^m$, hence is independent of the λ_i 's (i.e., the m -osculating spaces are constant along the generators of X).

Moreover, these spaces are the ones spanned by the $[D_j]$'s, $1 \leq j \leq r-1$, and the m -osculating spaces to D_r . In other words, in this case

$$X_m^\vee = S^{d_r-m-1}(D_r^*) \subset [X_1, \dots, X_{r-1}]^\vee$$

is the (d_r-m-1) -th osculating developable of the strict dual curve $D_r^* \subset [X_1, \dots, X_{r-1}]^\vee$. In particular $\dim X_m^\vee = d_r-m$.

Proof of Proposition 2:

Case 1): $d_{r-1} = d_r = d$.

Considering the matrix A_m we observe that $\bar{m} = d$ holds, and that every d -osculating space contains the space $[D_1, \dots, D_i] = [X_1, \dots, X_i]$. Moreover, for each k , $i+1 \leq k \leq r$, we have

$$X^*|_{D_k} = (D_{i+1}^*, \dots, D_{k-1}^*, D_{k+1}^*, \dots, D_r^*) \subset [X_1, \dots, X_i]^\vee \cap [D_k]^\vee.$$

Thus we see that

$$X^* = X_d^\vee = (D_{i+1}^*, \dots, D_r^*) \subset [X_1, \dots, X_i]^\vee$$

holds.

Case 2): $d_{r-1}+1 = d_r$.

We see that $\bar{m} = d_{r-1} = d$ holds, and that every d -osculating space contains $[X_1, \dots, X_i]$. Moreover, for each k , $i+1 \leq k \leq r-1$, we have

$$X^*|_{D_k} = (D_{i+1}^*, \dots, D_{k-1}^*, D_{k+1}^*, \dots, D_{r-1}^*, S^1(D_r^*)) \subset [X_1, \dots, X_i]^\vee \cap [D_k]^\vee,$$

where $S^1(D_r^*) \subset [X_1, \dots, X_{r-1}]^\vee$ is the tangent developable of D_r^* .

For $k = r$, we have

$$X^*|_{D_r} = (D_{i+1}^*, \dots, D_r^*) \subset [X_1, \dots, X_i]^\vee \cap [D_r]^\vee.$$

It follows that

$$X^* = X_d^\vee = (D_{i+1}^*, \dots, D_{r-1}^*, S^1(D_r^*)) \subset [X_1, \dots, X_i]^\vee.$$

To see that X^* is (non-normal) of type (d, \dots, d) , it suffices to show that $S^1(D_r^*)$ is (non-normal) of type (d, d) .

This follows from the following general lemma.

LEMMA: Let $D \subset \mathbb{P}^d$ be a rational normal curve of degree d ($d \geq 3$). Then the m -osculating developable $S^m(D)$ ($m \leq d-2$) of D is a (non-normal) variety of type $(d-m, \dots, d-m)$ (in \mathbb{P}^d). In other words, the bundle $P_D^m(1)$ is isomorphic to $\bigoplus_{i=0}^m \mathcal{O}_{\mathbb{P}^1}(d-m)$, i.e., $P_D^m(1)$ is semistable.

Proof: It suffices to exhibit $m+1$ independent sections of degree $d-m$. If D is given by

$$x_i = u^i v^{d-i}, \quad i = 0, 1, \dots, d,$$

then

$$s_k = \frac{\partial^m x}{\partial u^{m-k} \partial v^k}, \quad k = 0, \dots, m,$$

are the desired sections.

Case 3): This has already been shown in the proof of Prop. 1, by observing $\bar{m} = d_{r-1}$.

REMARK: Suppose $d_1 > 0$ (i.e. X is a rational normal scroll). Then biduality ($X^{**} = X$) holds if and only if $d_1 = \dots = d_{r-1}$ and $d_r = d_{r-1}$ or $d_r = d_{r-1} + 1$, in other words, if and only if X is a general scroll (in the sense of Ghione [G]). In the balanced case ($d_1 = \dots = d_r$) we have already seen that X is in fact self-dual. In the second case we have

$$X^{**}|_{D_j^*} = X_{i_1, \dots, i_{r-1}} \quad (j \neq r)$$

and

$$X^{**}|_{S^1(D_r^*)} = X_{1, \dots, r-1},$$

from which we conclude: $X^{**} = X$.

In all other cases, there is an i , $1 \leq i \leq r-1$, such that $d_{i+1} < \bar{m} < d_{i+1}$; hence

$$X^* \subset [X_1, \dots, i]^V.$$

Consequently, X^{**} is the cone over X with vertex $[X_1, \dots, i]$.

6. Curves in Grassmann varieties

A ruled variety can also be considered as a variety in a Grassmann variety. If $X \subset \mathbb{P}^N$ is of type (d_1, \dots, d_r) , defined by

$$V_{\mathbb{P}^1} \rightarrow \mathbb{P} = \bigoplus_{i=1}^r \mathcal{O}_{\mathbb{P}^1}(d_i), \text{ then that also defines a morphism}$$

$$\mathbb{P}^1 \rightarrow G = \text{Grass}_r(V).$$

Let $\Gamma \subset G$ denote the image of this map.

The m -dual variety $\Gamma_m^V \subset G^V = \text{Grass}_r(V^V)$ is defined as (the closure of) the set of $(N-r)$ -spaces that contain an m -osculating space to Γ [P2, §2]. These latter spaces are defined by the image of

$$b^m: V_{\mathbb{P}^1} \rightarrow \mathbb{P}_{\mathbb{P}^1}^m(\mathbb{P}) = \bigoplus_{i=1}^r \mathbb{P}_{\mathbb{P}^1}^m(d_i).$$

The image of b^m is defined by the matrix

$$B_m = \begin{pmatrix} \bar{M}_{d_1}^m & 0 \\ 0 & \bar{M}_{d_r}^m \end{pmatrix}$$

of rank $\beta_m = r(m+1) - \sum_{j=1}^i (m-d_j)$, where $i = i(m)$ is the integer such that $0 \leq i \leq r-1$ and $d_i \leq m < d_{i+1}$ (again we set $d_0=0$). The m -dual variety Γ_m^V is defined whenever $\beta_m \leq N+1-r$.

The strict dual variety of Γ is $\Gamma^* = \Gamma_{\bar{m}}^V \subset G^V = \text{Grass}_r(V^V)$, where \bar{m} is the smallest integer m such that β_m is maximal subject to $\beta_m \leq N+1-r$.

If $d_1 > 0$ and $m \leq d_1 - 1$, then $\beta_m = r(m+1)$ and Γ_m^V is defined. In this case, $\text{Ker}(b^m)^V$ is the sum of the bundles of principal parts of order $d_i - 1 - m$ of the "dual" of $\mathcal{O}_{\mathbb{P}^1}(d_i)$ (interpreted in terms of the curve D_i and its strict dual D_i^* , using duality for the corresponding short exact sequences ([P1], 5.2)).

PROPOSITION 3: Let $\Gamma \subset G$ correspond to $X \subset \mathbb{P}^N$ of type (d_1, \dots, d_r) , $r \geq 2$.

- 1) If $d_1 = \dots = d_r = d$, then $\Gamma^* = \Gamma_{d-1}^V$, $\dim \Gamma^* = 1$, and $X^* = X_d^V$ is the total space of Γ^* .
- 2) If $d_1 = \dots = d_{r-1} = d$, $d_r = d+1$, then $\Gamma^* = \Gamma_{d-1}^V$, $\dim \Gamma^* = 2$, and $X^* = X_d^V$ is the total space of Γ^* .
- 3) If $d_1 < d_2 = \dots = d_{r-1} = d$, $d_r = d+1$, then $\Gamma^* = \Gamma_{d-1}^V$, $\dim \Gamma^* = 1$.

In all other cases, $\Gamma^* = \Gamma_{\bar{m}}^V$, where $\bar{m} < d_{r-2}$ if $r \geq 3$; if $r = 2$, $d_{r-1} < d_{r-2}$, then $\bar{m} = d_{r-2}$, and Γ^* is the curve corresponding to the tangent developable of D_2^* .

Proof: Assume $d_r = d$, and that Γ_{d-1}^V is defined. Then (with i as before),

$$\beta_{d-1} = rd - \sum_{j=1}^i (d-1-d_j) \leq N+1-r = \sum_{j=1}^r d_j,$$

so

$$rd - i(d-1) \leq \sum_{j=i+1}^r d_j \leq (r-i)d.$$

Hence $i = 0$, and $d-1 < d_1 \leq d_2 \leq \dots \leq d$. Therefore $d_1 = \dots = d_r = d$.

In this case, clearly $\bar{m} = d-1$, and Γ^* is the curve whose total space is $X^* = (D_1^*, \dots, D_r^*)$.

Assume $d_{r-1} = d$, $d_r = d+1$, and that Γ_{d-1}^V is defined (clearly $\bar{m} \leq d-1$). Then

$$\beta_{d-1} = rd - \sum_{j=1}^i (d-1-d_j) \leq \sum_{j=1}^r d_j,$$

so that we get

$$rd - i(d-1) \leq \sum_{j=i+1}^r d_j \leq (r-i)d+1.$$

Hence $i \leq 1$. Suppose $i = 0$. Then $d-1 < d_1$, so $d_1 = \dots = d_{r-1} = d$, $d_r = d+1$, $\beta_{d-1} = rd$, and $\text{Ker}(b^{d-1})$ has rank

$$N+1-rd = (r-1)d+d+1+r-rd = r+1.$$

Hence $\Gamma^* = \Gamma_{d-1}^V$ is ruled by lines, so $\dim \Gamma^* = 2$.

The matrix B_{d-1} in this case is just the one associated with $X^* = X_d^V = (D_1^*, \dots, D_r^*, S^1(D_r^*))$.

If $i = 1$ (hence $r \geq 3$), then $\beta_{d-1} = (r-1)d+d_1+1$ and the rank of $\text{Ker}(b^{d-1})$ is $N+1-\beta_{d-1} = r$. Hence $\Gamma^* = \Gamma_{d-1}^V$ is a curve.

Suppose finally that $d_{r-1} < d_{r-2}$, and that Γ_m^V is defined for some $m \geq d_{r-2}$. Then $i = r-1$, and we must have

$$r(m+1)-(r-1) \leq d_r, \quad r+m \leq d_r \leq m+2.$$

Hence $r \leq 2$, so $r = 2$, and $m = d_2-2$. Then clearly Γ^* is the curve in G^V corresponding to the tangent developable $S^1(D_2^*)$ of D_2^* in $[D_2^*]$.

REMARK: Set $F_m^* = (\text{Ker } b^m)^V$. From what we have seen, F_m^* is the direct sum of bundles of principal parts on the curves D_1^*, \dots, D_r^* . We can write

$$F_m^* = \bigoplus_{j=1}^r \mathcal{P}_{\mathbb{P}^1}^{d_j-m-1}(d_j)$$

(if $m+1 > d_j$, we set the bundle equal to 0).

From a previous lemma, we know that $\mathcal{P}_{\mathbb{P}^1}^{d_j-m-1}(d_j)$ is balanced: it splits into d_j-m copies of $\mathcal{O}_{\mathbb{P}^1}(d_j-(d_j-m-1)) = \mathcal{O}_{\mathbb{P}^1}(m+1)$. Hence, for m such that $m \leq d_1-1$, the $(\sum_{j=1}^r d_j-rm)$ -bundle F_m^* is balanced of type $(m+1, \dots, m+1)$. Hence Γ_m^V is defined by a balanced bundle F_m^* (recall that Γ_m^V is the image of $\text{Grass}_r(F_m^*) \subset \mathbb{P}^1 \times \text{Grass}_r(V^V)$ in $G^V = \text{Grass}_r(V^V)$), even though the bundle F defining Γ was not necessarily balanced. Moreover, the case F balanced corresponds to F_{d-1}^* balanced and $\text{rank } F_{d-1}^* = r$; this is precisely when X and Γ are self-dual.

References

- [G] F. GHIONE, "Quot schemes over a smooth curve", Preprint, 1982.
- [P1] R. PIENE, "Numerical characters of a curve in projective n-space". In Real and complex singularities, Oslo 1976.
Ed. P. Holm, Sijthoff and Noordhoff, Groningen 1978, 475-495.
- [P2] R. PIENE, "A note on higher order dual varieties, with an application to scrolls". To appear in Proc.Symp.Pure Math.: Singularities, Arcata 1981.
- [S1] C. SEGRE, "Sulle rigate razionali in uno spazio lineare qualunque", Atti della R. Acc. delle scienze di Torino XIX, 265-282 (1883-84).
- [S2] C. SEGRE, "Sulle varietà a tre dimensioni composte di serie semplici razionali di piani", Atti della R.Acc. delle scienze di Torino XXI, 95-115 (1885-86).

Ragni Piene
University of Oslo
Institute of Mathematics
P.B. 1953, Blindern
Oslo 3
Norway

Gianni Sacchiero
Istituto Matematico
Università di Ferrara
Via Machiavelli, 35
44100 Ferrara
Italy